

Bending of a light ray within a dispersive medium in a uniform gravitational field

Bernard Linet *

Laboratoire de Mathématiques et Physique Théorique

CNRS/UMR 6083, Fédération Denis Poisson

Université François Rabelais, 37200 TOURS, France

Abstract

Recently, Dressel *et al* determined the vertical deflection of a light ray in a medium, with strong frequency-dependent dispersion, at rest in a uniform gravitational field. We take up this question within the general relativistic theory of the propagation of light in a dispersive medium, due to Synge.

In a recent paper [1], Dressel *et al* determined the vertical deflection Δz of a light ray in a medium, with strong frequency-dependent dispersion, at rest in a uniform gravitational field. Following their method, they found

$$\Delta z \approx -\frac{gL^2v_p}{2c^2v_g}, \quad (1)$$

where g is the acceleration due to the Earth gravity, L the covered distance and v_p and v_g are respectively the phase velocity and the group velocity of this dispersive medium.

We take up this question in a more general approach. We consider the general relativistic theory of the propagation of light in a dispersive medium, proposed by Synge [2] a long time ago. We first summarize briefly this theory in a curved space-time. Then, we carry out the exact calculations in the case of the Rindler space-time representing a uniform gravitational field. For a weak acceleration of gravity, we find again formula (1). In our knowledge, this approach of the problem has not been already performed.

We consider a space-time which is described by a metric $g_{\mu\nu}$ in a coordinate system (x^μ) . We adopt the signature $(-+++)$ for the metric. We use units in which $c = 1$. The dispersive medium is described by its 4-velocity V^μ , satisfying $g_{\mu\nu}V^\mu V^\nu = -1$. We now consider the propagation of a wave in a dispersive medium. It is defined by the equation $S(x^\mu) = \text{constant}$ where S is a function. The vector of propagation k^μ is defined by

$$k_\mu = \partial_\mu S. \quad (2)$$

*E-mail: linet@lmpt.univ-tours.fr

The frequency ω of the wave, relative to V^μ , has the expression

$$\omega = -k_\mu V^\mu. \quad (3)$$

The phase velocity v_p , relative to V^μ , is given by

$$\frac{1}{v_p^2} = 1 + \frac{g^{\mu\nu} k_\mu k_\nu}{(k_\mu V^\mu)^2}. \quad (4)$$

In classical theory of optics, a dispersive medium is characterized by the refractive index n which is the reciprocal of the phase velocity, i.e.

$$n = \frac{1}{v_p}, \quad (5)$$

and consequently the relation of dispersion takes the form

$$g^{\mu\nu} k_\mu k_\nu - (n^2 - 1) \omega^2 = 0. \quad (6)$$

The refractive index n is a given function $n(x^\mu, \omega)$ which characterizes the dispersive medium.

The equations of the light rays are derived from the Hamiltonian function

$$\mathcal{H}(x^\mu, k_\nu) = \frac{1}{2} \left[g^{\mu\nu}(x^\lambda) k_\mu k_\nu - (n^2(x^\lambda, \omega) - 1) \omega^2 \right] \quad (7)$$

where $\omega = \omega(x^\lambda, k_\sigma)$ given by (3). The characteristic curves of $\mathcal{H}(x^\mu, k_\nu) = 0$ are the light rays. From the usual canonical equations, we find thus the following system of differential equations for a parameter λ :

$$\frac{dx^\mu}{d\lambda} = g^{\mu\nu} k_\nu + \left[(n^2 - 1) \omega + n \frac{\partial n}{\partial \omega} \omega^2 \right] V^\mu, \quad (8)$$

$$\frac{dk_\mu}{d\lambda} = -\frac{1}{2} \partial_\mu g^{\alpha\beta} k_\alpha k_\beta + n \partial_\mu n \omega^2 - \left[(n^2 - 1) \omega + n \frac{\partial n}{\partial \omega} \omega^2 \right] k_\nu \partial_\mu V^\nu, \quad (9)$$

and we have to add constraint (6).

The curves $x^\mu(\lambda)$ define the congruence of world-lines of light rays, i.e. the trajectories of the pulses of light. So, the vector $dx^\mu/d\lambda$ must be timelike or null. The group velocity v_g , relative to V^μ , is given by

$$v_g^2 = 1 + \frac{g_{\mu\nu} \frac{dx^\mu}{d\lambda} \frac{dx^\nu}{d\lambda}}{\left(\frac{dx^\mu}{d\lambda} V_\mu \right)^2}. \quad (10)$$

According to (8), we easily see that

$$g_{\mu\nu} \frac{dx^\mu}{d\lambda} \frac{dx^\nu}{d\lambda} = n^2 \omega^2 \left[1 - \left(\frac{\partial}{\partial \omega} (n\omega) \right)^2 \right] \quad \text{and} \quad \frac{dx^\mu}{d\lambda} V_\mu = -n\omega \frac{\partial}{\partial \omega} (n\omega). \quad (11)$$

By inserting expressions (11) into (10), we obtain ¹

$$v_g = \frac{1}{\frac{\partial}{\partial \omega}(n\omega)}. \quad (12)$$

We assume that the solutions to differential equations (8) and (9) yield $v_g \leq 1$.

We add the following property in the case where there exists a vector ξ^μ which is a Killing vector for the metric and also for the dispersive medium, i.e.

$$\mathcal{L}_\xi g_{\mu\nu} = 0 \quad , \quad \mathcal{L}_\xi n = 0 \quad \text{and} \quad \mathcal{L}_\xi V^\mu = 0 \quad (13)$$

where $\mathcal{L}_\xi$ is the Lie derivative with respect to ξ^μ . From equations (8) and (9), we can straightforwardly deduce the following differential equation along the light rays:

$$\frac{d}{d\lambda} (\xi^\mu k_\mu) = 0, \quad (14)$$

and hence we get that $\xi^\mu k_\mu$ is a constant of motion.

We are now in a position to consider the case of the Rindler space-time described by the metric

$$ds^2 = -(1 + gz)^2 (dx^0)^2 + dx^2 + dy^2 + dz^2, \quad (15)$$

well defined for $gz > -1$. When $|gz| \ll 1$, metric (15) represents a Newtonian potential $U = -gz$ in Galilean coordinates where g is the acceleration of the gravity. We assume that the dispersive medium is at rest in these coordinates and moreover that the refractive index n does not depend on the position, $n = n(\omega)$. The 4-velocity V^μ has the following components:

$$V^0 = \frac{1}{1 + gz} \quad \text{and} \quad V^x = V^y = V^z = 0. \quad (16)$$

In metric (15) with 4-velocity (16), there are three Killing vectors $\partial/\partial x^0$, $\partial/\partial x$ and $\partial/\partial y$.

We consider a light ray in the plane $x = 0$, horizontally emitted at the point $y = 0$ and $z = 0$. The constants of motion (14) are denoted

$$k_0 = -\omega_0 \quad , \quad k_x = 0 \quad \text{and} \quad k_y = \omega_0 k \quad (17)$$

where ω_0 and k are constants. According to (3), we notice that

$$\omega = \frac{\omega_0}{1 + gz}. \quad (18)$$

We see from (18) that ω_0 is the frequency of the wave, relative to V^μ , at $z = 0$. We write down equation (8) in this particular case

$$\frac{dx}{d\lambda} = 0 \quad , \quad \frac{dy}{d\lambda} = \omega_0 k \quad \text{and} \quad \frac{dz}{d\lambda} = k_z. \quad (19)$$

¹As noticed by Bičák and Hadrava [3], there is a misprint in the book of Synge [2].

We now give the relation of dispersion (6) with expressions (17) and (18)

$$\omega_0^2 k^2 + (k_z)^2 - n^2(\omega) \frac{\omega_0^2}{(1 + gz)^2} = 0. \quad (20)$$

At the point $y = 0$ and $z = 0$, we have $k_z = 0$ since the light is horizontally emitted. Consequently, we get from relation (20) at $z = 0$ the value of the constant k

$$k = n(\omega_0). \quad (21)$$

Finally, we obtain the desired expression of k_z

$$(k_z)^2 = \frac{n^2(\omega)\omega_0^2}{(1 + gz)^2} - n^2(\omega_0)\omega_0^2. \quad (22)$$

Taking into account (19), we find from (22) the exact differential equation

$$\left(\frac{dz}{dy}\right)^2 = \frac{n^2\left(\frac{\omega_0}{1 + gz}\right)}{(1 + gz)^2 n^2(\omega_0)} - 1 \quad (23)$$

giving the vertical deflection of the light ray when the refractive index $n(\omega)$ of the dispersive medium is known.

We are now ready to determine the vertical deflection in the limit where $|gz| \ll 1$. We expand expression (23) linearly in gz

$$\left(\frac{dz}{dy}\right)^2 \approx -\frac{2\frac{dn}{d\omega}(\omega_0)\omega_0 gz}{n(\omega_0)} - 2gz. \quad (24)$$

Making use of (4) and (12) expressed at $z = 0$, we rewrite (24) under the form

$$\left(\frac{dz}{dy}\right)^2 \approx -2\left(\frac{v_p}{v_g}\right)gz, \quad (25)$$

with $z < 0$ of course. Differential equation (25) leads obviously to formula (1) of Dressel *et al* [1].

We have obtained a general result. We have not made the slow light approximation that the refractive index is linear in the frequency.

References

- [1] J. Dressel, S. G. Rajeev, J. C. Howell and A. N. Jordan, Phys. Rev. A **79** 013834 (2009).
- [2] J. L. Synge, *Relativity: The General Theory*, Chap. XI, North-Holland Publishing Company (1966).
- [3] J. Bičák and P. Hadrava, Astron. Astrophys. **44** 389 (1975).